



Low back pain: critical assessment of various scales

Amit Garg¹ · Hardik Pathak¹ · Maxim V. Churyukanov^{2,3} · Rajendra B. Uppin⁴ · Tatyana M. Slobodin⁵

Received: 31 July 2019 / Revised: 2 December 2019 / Accepted: 29 December 2019 / Published online: 8 January 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Abstract

Purpose To study the various pain assessment tools based on their psychometric properties and ease of use.

Methods Published articles on psychometric properties of pain tools were accessed and data collected for low back pain (LBP)-specific tools, generic tools, neuropathic LBP tools, tools for cognitively impaired patients, and tools for acute LBP.

Results Among the LBP-specific tools, Roland Morris Disability Questionnaire (RMDQ) and Oswestry Disability Index (ODI) have good construct validity and reliability, and responsiveness over short intervals. Quebec Back Pain Disability Scale (QBPDS) gauges only disability and sleep. Among the generic tools, McGill Pain Questionnaire (MPQ), West Haven-Yale Multidimensional Pain Inventory (MPI), and Brief Pain Inventory (BPI) show good responsiveness, but BPI is the only tool validated for LBP. Neuropathic Pain Scale (NPS) and Short Form-MPQ-2 (SF-MPQ-2) are both reliable tools for neuropathic LBP. For cognitively impaired patients, Pain Assessment in Advanced Dementia (PAINAD), Abbey Pain Scale (APS), and Doloplus-2 are all reliable tools, but PAINAD has good construct validity. For acute pain, Clinically Aligned Pain Assessment (CAPA) is reliable and responsive, but presently, unidimensional tools and SF-MPQ-2 are the tools most preferred.

Conclusion Based on psychometric properties and ease of use, the best tools for LBP seem to be RMDQ/ODI (among LBP-specific tools), BPI (among generic tools), SF-MPQ-2/NPS (for neuropathic LBP), PAINAD (for cognitively impaired patients), and unidimensional tools and SF-MPQ-2 (for acute pain). Overall, BPI seems to be a tool that can be relied upon the most.

Graphic abstract

These slides can be retrieved under Electronic Supplementary Material.

The graphic abstract consists of three slides from a presentation. The first slide, titled 'Key points', lists three main findings: 1) Measuring low back pain is challenging due to patient description accuracy. 2) Simple pain assessment tools are needed for elderly and cognitively-impaired individuals. 3) Major pain assessment tools were critically assessed using published data. The second slide is a flowchart titled 'Major Pain Assessment Tools' which categorizes tools into 'LBP-Specific Tools' (RMDQ, ODI, QBPDS) and 'Generic Tools' (BPI, SF-MPQ-2, NPS, MPI, MPQ). The third slide, 'Take Home Messages', summarizes the best tools for different patient groups: RMDQ/ODI for LBP, BPI for generic use, SF-MPQ-2/NPS for neuropathic LBP, PAINAD/APS/Doloplus-2 for cognitively-impaired patients, and unidimensional tools/SF-MPQ-2 for acute pain.

Keywords Low back pain tools · Psychometric properties · RMDQ · ODI · BPI

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00586-019-06279-5>) contains supplementary material, which is available to authorized users.

✉ Hardik Pathak
drhardikpathak@gmail.com

¹ Global Medical Affairs, Dr. Reddy's Laboratories Ltd, Hyderabad, India

² The I.M. Sechenov First Moscow State Medical University (Sechenov University), Moscow, Russia

³ The B.V. Petrovsky Russian Scientific Surgery Center, Moscow, Russia

⁴ Department of Orthopaedics, KLE Academy of Higher Education and Research, JN Medical College, Belagavi, India

⁵ Shupyk National Medical Academy of Postgraduate Education, Kiev, Ukraine

Abbreviations

APS	Abbey Pain Scale
BPI	Brief Pain Inventory
CAPA	Clinically Aligned Pain Assessment
Doloplus	After the French investigator who developed the tool
FPS	Faces Pain Scale
MPI	West Haven-Yale Multidimensional Pain Inventory
MPQ	McGill Pain Questionnaire
MVAS	Million's Visual Analogue Scale
NASS-LSO	The North American Spine Society Lumbar Spine Outcome
NPS	Neuropathic Pain Scale
ODI	Oswestry Disability Index
NRS	Numerical Rating Scale
PAINAD	Pain Assessment in Advanced Dementia
QBPDS	Quebec Back Pain Disability Scale
RMDQ	Roland Morris Disability Questionnaire
SF-36	Short Form-36
VAS/GRS	Visual Analogue Scale/Graphic Rating Scale
VRS	Verbal Rating Scale
WDI	Waddell Disability Index

Introduction

Pain afflicts humans not only in sickness, critical illness, or palliative care but also in their everyday life. According to International Classification of Functioning, Disability and Health, low back pain (LBP) alters basic body functions such as sleep and motility, and causes changes in a person's spinal and musculoskeletal structures [1]. Acute and chronic LBP can be caused by a variety of medical conditions including trauma, muscle strain, lumbar spondylosis, spinal stenosis, spine and kidney infections, certain cancers, endometriosis, arthritis, and ankylosing spondylitis. More than 80% of the population experiences LBP at some point in their lives [2]. Many go on to recover naturally or with the help of some form of intervention, while others develop chronic back pain with pain lasting more than 3 months. The number of years lived with disability triggered by LBP has increased by 54% between 1990 and 2015, worldwide [3]. LBP is one of the major reasons of low performance at work, missed work, disability requests after an episode of acute pain or post-operative condition, and is known to cause more global disability than any other condition [4–6].

The major domains assessed for chronic pain are pain severity, pain-related functional interference, and emotional burden [7], while for acute pain it is mainly pain severity and relief thereof [8]. Pain assessment tools can

be used for general pain (generic tools), for condition-specific pain, or for health-related quality of life (HRQoL) [9]. Many general tools are used independently in disease conditions or used in combination with some condition-specific tool. A condition-specific tool is also used in conjunction with some HRQoL tool for complete pain assessment. Tools can be self-reported or observational depending upon the patient's consciousness or cognizance. In this article, we aim to discuss some major generic, condition-specific, and HRQoL tools that are used regularly and extensively by physicians and patients worldwide for LBP.

Search strategy

For this review, we first set a few primary questions:

1. What are the tools/scales available for the assessment of chronic and acute pain?
2. What tools are majorly used for chronic and acute LBP?
3. How are these tools relevant for various patient populations like people with and without cognitive impairment, and neuropathic LBP?

Search terms were entered as Medical Subject Headings (MeSH) terms or free text on PubMed and Scopus. Following terms were used to search literature: “pain assessment scales”, “chronic back pain”, “low back pain assessment”, “psychometric properties”, “patient-reported outcome measures”, “PROM”, “dementia”, “neuropathic pain”, “pain in elderly”. Original research articles and reviews (systematic and narrative) from January 1980 to February 2019 were accessed. Articles without a comprehensive study of pain assessment tools were excluded. Articles on pain assessment tools, specifically for psychometric properties, were searched to understand what the major patient-reported outcome measures (PROM) were. Comprehensive studies and reviews on the validity, reliability, stability, and responsiveness of each PROM were selected. We found that certain generic tools [Brief Pain Inventory (BPI), McGill Pain Questionnaire (MPQ), West Haven-Yale Multidimensional Pain Inventory (MPI), and Short Form-36 (SF-36)] are extensively used to assess LBP, and thus, we extended our search to include these tools along with LBP-specific tools. These generic tools are used to measure pain in general that may not be localized to a specific region of the body. A list was drawn for the number of articles obtained for nine major tools suggested [9, 10]. Among these, there were numerous results (> 100) for three tools [Roland Morris Disability Questionnaire (RMDQ), Oswestry Disability Index (ODI),

Table 1 Selection of low back pain-specific tools

S. no.	Name of low back pain-specific tool	Review articles + systematic reviews + RCTs	Foreign language + irrelevant topics (not LBP + does not involve the pain tool)	Relevant articles
1.	ALBPS	28	11	17
2.	LBOS	9	3	6
3.	LBPRS	9	4	5
4.	MVAS	78	46	32
5.	NASS	49	3	46
6.	ODI	3678	NA	3678
7.	QBPDS	159	NA	159
8.	RMDQ	976	NA	976
9.	WDI	36	10	26

ALBPS Aberdeen LBP Scale, *LBOS* Low Back pain Outcome Score, *LBPRS* Low Back Pain Rating Scale, *MVAS* Million's Visual Analogue Scale, *NASS-LSO* The North American Spine Society Lumbar Spine Outcome, *ODI* Oswestry Disability Index, *QBPDS* Quebec Back Pain Disability Scale, *RMDQ* Roland Morris Disability Questionnaire, *WDI* Waddell Disability Index

Table 2 Pain assessment tools based on various outcome domains and patient groups

S. no.	Outcome domains	Name of tools
1.	Pain intensity, quality, location, and pattern (general and condition-specific)	MPQ, RPS
2.	Pain interference and functioning (general)	PDI, BPI, FIM, DDS
3.	Pain interference and functioning (disease-specific)	WOMAC, RMDQ, ODI, QBPDS, NASS-LSO, MVAS, WDI, MPI, SF-36, SIP, NPS, PAINAD, APS, Doloplus-2
4.	Psychosocial measurements	BDI, PCS, CSQ
5.	Observational pain assessment instruments (general and critically ill patients). Mostly for non-verbal pain assessment when patient may not be alert, conscious or in a state to talk	CPOT, ESAS, PCOS

APS Abbey Pain Scale, *BDI* Beck Depression Inventory, *BPI* Brief Pain Inventory, *CSQ* Coping Strategies Questionnaire, *CPOT* Critical Care Pain Observation Tool, *DDS* Descriptor Differential Scale, *Doloplus-2* After the tool's developer, *ESAS* Edmonton Symptom Assessment Scale, *FIM* Functional Independence Measure, *MPQ* McGill Pain Questionnaire, *MVAS* Million Visual Analogue Scale, *NPS* Neuropathic Pain Scale, *ODI* Oswestry Disability Index, *PAINAD* Pain Assessment in Advanced Dementia Scale, *PCS* Pain Catastrophizing Scale, *PDI* Pain Disability Index, *PCOS* Palliative Care Outcome Scale, *QBPDS* Quebec Back Pain Disability Scale, *RPS* Regional Pain Scale, *RMDQ* Roland Morris Disability Questionnaire, *SF-36* Short-Form survey, *SIP* Sickness Impact Profile, *NASS-LSO* The North American Spine Society Lumbar Spine Outcome, *WDI* Waddell Disability Index, *MPI* West Haven-Yale Multidimensional Pain Inventory, *WOMAC* Western Ontario McMaster Osteoarthritis Index

Quebec Back Pain Disability Scale (QBPDS)], and their popularity evident, we did not go through all the results to check for data relevance. Three other tools [Aberdeen LBP Scale (ALBPS), Low Back pain Outcome Score (LBOS), and Low Back Pain Rating Scale (LBPRS)] were excluded from this review due to lack of substantial results, bringing it to a total of six LBP-specific tools [RMDQ, ODI, QBPDS, Million Visual Analogue Scale (MVAS), Waddell Disability Index (WDI), and the North American Spine Society Lumbar Spine Outcome (NASS-LSO)] (Table 1).

Tools/scales for measurement of chronic and acute pain

In 2002, Initiative on Methods, Measurement, and Pain Assessment in Clinical Trials (IMMPACT) was developed by a team of pain experts from various industries and academia in the USA as an improved method of pain assessment during clinical trials [11]. They highlighted the importance of six core outcome domains to assess chronic pain: pain, physical functioning, emotional functioning, patient's rating of overall improvement after treatment, patient-reported adverse effects during treatment, and patient's disposition and characteristics data. In the same year, more outcome domains pertaining to HRQoL were also added to IMMPACT [12]. There are various pain assessment tools




Fig. 1 Majorly used pain assessment tools. *APS* Abbey Pain Scale, *BPI* Brief Pain Inventory, *CAPA* Clinically Aligned Pain Assessment, *Doloplus* After the French investigator who developed the tool, *FPS* Faces Pain Scale, *MPI* West Haven-Yale Multidimensional Pain Inventory, *MPQ* McGill Pain Questionnaire, *MVAS* Million’s Visual Analogue Scale, *NASS-LSO* The North American Spine Society

Lumbar Spine Outcome, *NPS* Neuropathic Pain Scale, *ODI* Oswestry Disability Index, *NRS* Numerical Rating Scale, *PAINAD* Pain Assessment in Advanced Dementia, *QBPDS* Quebec Back Pain Disability Scale, *RMDQ* Roland Morris Disability Questionnaire, *SF-36* Short Form-36, *VAS/GRS* Visual Analogue Scale/Graphic Rating Scale, *VRS* Verbal Rating Scale, *WDI* Waddell Disability Index

Table 3 Unidimensional pain assessment tools

S. no.	Name of tool	Description
1.	NRS	A 0–10/0–20/0–100 scale of pain intensity, where zero and maximum value stand for no pain and highly intense pain, respectively
2.	VAS/GRS	Patients select their level of pain on a 10 cm scale; sometimes with descriptive terms
3.	VRS	4–5-point scale with words like “no pain”, “moderate pain”, “severe pain”, etc. SF-36 is a good example
4.	FPS	Mostly for children and elderly, choices range between a happy face to a face showing extreme pain

FPS Faces Pain Scale, *NRS* Numerical Rating Scale, *SF-36* Short Form-36, *VAS/GRS* Visual Analogue Scale/Graphic Rating Scale, *VRS* Verbal Rating Scale

associated with different outcome domains and patient groups (Table 2) (Fig. 1).

Chronic low back pain tools

For the assessment of chronic LBP, both unidimensional and multidimensional tools are used.

Unidimensional tools These tools only measure the intensity of pain and are explained briefly in Table 3.

Multidimensional tools These tools measure not only the severity of pain but also integrate other dimensions such as pain-based interference on HRQoL and functioning,

location of pain, and emotional burden. The majorly used multidimensional tools for adults (young and elderly) with and without cognitive impairment, and for adults with neuropathic low back pain are explained in Tables 4 and 5. The merits and demerits of these tools are listed in Table 6.

Acute low back pain tools

Acute pain measurement and management is usually straightforward and relies on documenting two outcome domains, namely pain intensity and relief post-treatment.– They usually involve unidimensional tools such

Table 4 Multidimensional tools for adults with and without cognitive impairment, and for adults with neuropathic low back pain

Multidimensional tools for adults (young and elderly) without cognitive impairment		
Generic tools	LBP-specific tools	
BPI [13]	<p>15-item tool</p> <p>2 domains: pain intensity, pain interference</p> <p>Short-form BPI for both chronic and acute pain LBP</p> <p>Valid and reliable in accurately assessing pain</p>	<p>RMDQ-24 [14]</p> <p>24 questions</p> <p>2 Psychosocial</p> <p>3 Sleep/rest</p> <p>15 Physical functioning</p> <p>1 Eating</p> <p>2 Home management</p> <p>1 back pain frequency</p> <p>5 min to complete</p> <p>Questions pertain to pain experienced during various activities in and around the house</p> <p>Most questions include the phrase “because of my back”</p> <p>Questions:</p> <p>RMDQ-23/RMDQ-18: derived versions</p>
MPQ [15]	<p>New 22-item version by Dworkin et al. [16] that includes neuropathic pain</p> <p>Useful for both chronic and acute pain</p> <p>SF-MPQ-2: 15 items derived version</p>	<p>ODI [17]</p> <p>10 items</p> <p>5 min to complete</p> <p>Includes questions on pain intensity, sex life, social life, and travelling</p>
MPI [18]	<p>52 items, 12 scales</p> <p>3 parts:</p> <p>Pain severity, pain interference, Responses</p> <p>Household, social activities</p>	<p>QBPDS [19]</p> <p>20 items</p> <p>6 major groups</p> <p>5–10 min to complete</p> <p>No questions on social life, sex life, and pain intensity</p>
SF-36 [20]	<p>36 items</p> <p>2 health constructs: PCS, MCS</p> <p>5–10 min to complete</p>	<p>MVAS [21]</p> <p>15 questions</p> <p>3 major domains</p> <p>Body functions</p> <p>Daily activities</p> <p>Social life</p> <p>5 min to complete</p> <p>Uses VAS</p> <p>WDI [22]</p> <p>9 items</p> <p>5 min to complete</p> <p>Items on back-involved activities and social life</p> <p>NASS-LSO [23]</p> <p>62 items</p> <p>5 categories</p> <p>Demographic data</p> <p>Medical history</p> <p>Body functions</p> <p>Employment history</p> <p>Treatment outcomes</p> <p>20 min to complete</p> <p>Pain locating picture to mark the regions of pain</p>
Multidimensional tools for cognitively impaired adults with low back pain		
PAINAD [24]	<p>5 items: facial expression, breathing, negative vocalizations, consolability, body language</p> <p>Range: 0–10</p> <p>Only few minutes to complete</p> <p>Endorsed by American Medical Directors Association</p> <p>Derived from items of FLACC, DS-DAT, and pain experiences of patients and clinical staff</p>	
APS [25]	<p>6 items: facial expression, vocalization, physiological change, physical change, change in body language, behavioural change</p> <p>Range: 0–3</p> <p>Less than a minute to complete</p> <p>Recommended by British Geriatrics Society and the Australian Pain Society</p>	
Doloplus-2 [26]	<p>10 items</p> <p>5 somatic, 2 psychomotor, 3 psychosocial items</p> <p>6–12 min to complete</p> <p>Overall score of 0–30</p>	

Table 4 (continued)

Multidimensional tools for adults with neuropathic low back pain

NPS [27]	10 items 8 items for pain localization and quality, 2 global items for pain severity and unpleasantness
----------	--

APS Abbey Pain Scale, BPI Brief Pain Inventory, *Doloplus* After the French investigator who developed the tool, FLACC Face, Legs, Activity, Cry, Consolability Scale, MPI West Haven-Yale Multidimensional Pain Inventory, MPQ McGill Pain Questionnaire, MVAS Million's Visual Analogue Scale, MCS Mental Component Summary, NASS-LSO The North American Spine Society Lumbar Spine Outcome, NPS Neuropathic Pain Scale, ODI Oswestry Disability Index, PCS Physical Component Summary, PAINAD Pain Assessment in Advanced Dementia, QBPDS Quebec Back Pain Disability Scale, RMDQ Roland Morris Disability Questionnaire, SF-36 Short Form-36, SF-MPQ-2 Short Form-MPQ-2, WDI Waddell Disability Index

as Numerical Rating Scale (NRS), Visual Analogue Scale (VAS), Verbal Rating Scale (VRS), and Faces Pain Scale (FPS), but multidimensional tools like BPI and SF-MPQ-2 are also used to assess pain. Acute pain, like chronic pain, should be assessed at both rest and movement, but movement-evoked pain assessment is more crucial for acute pain as this variety of pain is also a part of post-operative recuperative pain. There are certain unvalidated mnemonic tools as well, and even though they are unvalidated, they are very useful in measuring acute pain [8] (Table 7). Apart from these unvalidated tools, there is a new tool called Clinically Aligned Pain Assessment (CAPA) that is being used successfully for acute pain management. In 2013, Donaldson and Chapman [8, 64] developed this tool for University of Utah Health Care to replace the unidimensional NRS. This tool is now being used primarily in the University of Utah Hospital. A total of five domains, namely comfort, change in pain, pain control, functioning, and sleep are assessed. There is no script to follow or boxes to check, and patients and physicians engage in a conversation that is later evaluated in a coded manner. In a quality improvement study with more than 12,000 pain assessment observations, patients opted for CAPA much more than NRS [8]. Topham and Drew came up with its modified version in 2015 [64]. They used both process and outcome measures to understand staff compliance and patient satisfaction, respectively. This measure proved to be successful in increasing patient and staff satisfaction for both the independent research groups [8, 64]. Reliability and validity of the tool are not tested as it lacks scoring rules.

Comparing the different tools for assessment of low back pain

The biggest challenge faced by physicians and nursing staff is the ability to capture the severity of chronic pain as most often they have to rely on the patient's memory or description of pain that may not be accurate. When daily diaries of pain are to be maintained by the patient, they are usually

filled up in a hurry just prior to the visit to the doctor. Other limitations to capturing pain accurately are factors such as lack of education, disability, language barriers, incomprehension due to dementia, or simply, the lack of consciousness. Some pain scales can be complex and require training to use, while others can be long, making most respondents not answer all items in the questionnaire [26, 54]. Elderly and cognitively impaired individuals require pain scales that are simple, yet comprehensive [56, 57]. Thus arose the need to formulate specific scales that are simple, quick, and can be used to not only acquire the patient's input but also deliver quality pain management, providing eventual satisfaction and relief to both the patient and physician. Over the years, many tools have been developed for the assessment and management of LBP. Evaluating the quality of a tool to understand its psychometric properties is very important in helping the physician choose the best possible instrument to use [43]. In this review, we tried to maintain a level of consistency in reviewing all the psychometric properties of a tool, but there was a gap in information/research on certain parameters.

For the assessment of LBP, functional/disability measures, pain severity measures, and the measures for the effect of pain on social life and general mental health are used [9]. Functional measures can be both generic and condition-specific. Generic measures estimate symptoms related to various regions of the body and need not be spine-centric, while condition-specific measures assess symptoms specific to a disease or condition, for example, LBP. Both generic and condition-specific measures are used by physicians and patients as functional measures for LBP management [65, 66]. All LBP-specific measures that we covered in this article have been successfully used cross-culturally.

RMDQ and ODI have good content, construct validity, test-retest reliability, and responsiveness but are good at detecting changes over short intervals only [43]. While ODI can be used as both a functional measure and for pain severity, RMDQ lacks there in being only a good functional measure. Evaluation of psychological domains cannot be done with RMDQ as certain questions pertaining to the areas are lacking. It works well on patients with moderate

Table 5 Psychometric properties of low back pain tools (generic and condition-specific)^{a,c,e}

Pain assessment tool	Reliability	Validity	Responsiveness	Floor-ceiling effects
<i>Generic tools for back pain</i>				
BPI	High test–retest reliability value of 0.97 [28] Internal consistency for pain severity and interference: 0.89, 0.95 for arthritis, 0.82, 0.93 for low back pain [29]	High correlations with body pain scale of SF-36 (0.61–0.64), and RMDQ (0.57–0.81) [29]	Excellent sensitivity; SRM values for pain severity and interference – 1.09 and – 1.13, respectively [29]	Ceiling effect [30]: Subscale ‘pain 24-h worst’—18.5% Other subscales—1.4–3.5%
MPQ	Test–retest reliability values mostly above 0.9 [16, 31] Test–retest values of 0.62–0.75 (musculoskeletal pain) and 0.79–0.95 (rheumatic pain) [32] Internal consistency at 0.88–0.91 [31]	Poor to fair correlations of 0.21–0.42 with short-form BPI and NRS [30]	Sensitivity of the tool at 1–3 day interval is comparable with that of VAS ROC moderate at 0.61; SRM large at > 0.8 suggesting good sensitivity to change [32]	Floor effect: 12.5% and 15.1% for neuropathic and affective subscales, respectively [30]
MPI	Fair to excellent test–retest reliability (0.62–0.91; 0.46–0.93; 0.62–0.89), [18, 33, 34] Significant internal consistency values of 0.60–0.92 (cancer) [35], 0.63–0.93 (chronic back pain) [33]	Excellent content validity suggesting all items within the tool measure the same construct [35] Pain interference of MPI (0.76) relates well with BDI (0.51) [18]	Fair to moderate SRM between – 0.10 to 0.72 (various subscales) [34]	No floor or ceiling effects reported
<i>HRQL tool</i>				
SF-36	Good test–retest reliability values > 0.8 for PF, PR and BP, fair values for MH (0.55) [36] Good internal consistency ≥ 0.78 for all scores [37]	Good validity as all items are about health or pain but wide floor-ceiling effects of many items for rheumatic patients may be a sign that all health concerns are not fully captured [36] Correlation values of – 0.295 to – 0.66 for PF, BP and MH for RMDQ [38]; – 0.428 to – 0.70 for PF, BP, SF and MH for ODI [38]; – 0.25 to – 0.64 for PF, BP and MH for QBPDS [39]	SRM scores 0.44–0.67 [40]	Large floor-ceiling effects observed with rheumatic patients [36]
<i>Low back pain-specific tools</i>				
RMDQ-24	Test–retest reliability values 0.91 for same day [14], 0.86 for 3–6 weeks [41] Excellent internal consistency of 0.86 for 7 days [42]	Limited range of questions pertaining to physical functions; designed to address general problems faced by people suffering from back pain; only one question applies to the emotional state of the person [43] No certain content validity [44]; fair correlation of 0.5 [45] to ODI, and 0.6 to QBPDS [45]	Relevant for patients with little disability [43] Retest interval ROC values at 6 weeks (0.77) [40], 8 weeks (0.64) [46]	Floor effects present but no ceiling effects [47]
ODI	Test–retest reliability value for 1 day (0.94–0.99) [48, 49] Inter-rater reliability high (0.94) [49] Internal consistency values at 0.835–0.90 [49]	Uncertain content validity [44]. Correlation with RMDQ of 0.5 correlation coefficient [45]	High sensitivity with ROC values > 0.7 [40, 46]	Floor effects present but no ceiling effects [47]

Table 5 (continued)

Pain assessment tool	Reliability	Validity	Responsiveness	Floor-ceiling effects
QBPDS	Test–retest reliability values of 0.84 at 6 weeks, and 0.93 at 1–14 days [19, 39] Internal consistency > 0.9 [19]	Good content with all questions pertaining to various activities; agrees with both physicians and patients No questions on sexual activities Strong correlation with RMDQ (0.77), ODI (0.80), physical function subscale of SF-36 (0.72) [19]	Extremely sensitive tool with ROC values at 0.74 and above 0.80 [40]	No floor or ceiling effects [40]
MVAS	Test–retest reliability values 0.735–0.844 (daily activities), 0.581–0.745 (body functions), 0.736–0.849 (social life) [50] Internal consistency acceptable for daily activities and social life (0.794, 0.799) but unacceptable for body functions (0.67) [50]	Excellent goodness of fit results with confirmatory factor analysis; suggests all questions of the tool are essential and contribute to measuring what the tool sets out to assess [50]	Significant decrease in mean pain at three-month and one-year interval (17.1, 15.9) [51] Predicted treatment outcomes in patients with chronically disabling spine disorder [52]	Floor and ceiling effects below 15% [50]
WDI	Moderate test–retest reliability (ICC = 0.74) for “unchanged group” [40] Per cent inter-rater reliability 73–90% for each question of the tool [22]	Factor analysis confirmed the tool’s questions to be consistent, interconnected and to provide a good measure of disability [22] Correlates well with another disability tool, ODI, with correlation coefficient of $r=0.7$ [22]	Able to detect significant changes after one-month interval in patients [53]	No reported floor-ceiling effects
NASS-LSO	High test–retest reliability for 24 h (0.85–0.97) [23] High internal consistency (0.71–0.93) [23] Test–retest reliability 0.82 (pain/disability), 0.95 (neurogenic symptoms) and 0.63 (expectations) [54]	Pain/disability functions correlate well with SF-36 PCS (–0.82) and VAS (0.46–0.63) at baseline [54] Correlation with the MCS of SF-36 only fair with a value of 0.5 [54]	Large effect size (2.02) for combined pain/disability and neuro symptoms [54]	No reported floor-ceiling effects
<i>Tools for cognitively impaired</i>				
PAINAD	Good inter-rater reliability of 0.82–0.97 during pleasant and unpleasant activities [24], 0.31 at rest and 0.54 during movement [55] Internal consistency 0.50–0.6 [24]	Good construct; can discriminate between pain associated with different conditions, like pleasant and unpleasant activities High concurrent validity values of 0.89–0.95 (pleasant activity) and 0.82–0.91 (unpleasant activity) with pain scale of VAS [24] Good correlations (0.76) with DS-DAT and discomfort scale of VAS [24]	No reported measure of responsiveness	No reported floor-ceiling effects

Table 5 (continued)

Pain assessment tool	Reliability	Validity	Responsiveness	Floor-ceiling effects
APS	Inter-rater reliability values of 0.83 and 0.88 at rest and exercise, respectively [56]; internal consistency of 0.65–0.736 [57] Removing the item “physical change” improved the fit of the tool and increased internal consistency to 0.77 from 0.75 [56]	Correlation coefficient of APS with PAINAD 0.75–0.82 (at rest), 0.83–0.89 (during exercise), and 0.69–0.83 (at rest), 0.68–0.79 (during exercise) with DS-DAT [56]; indicates that all three tools measure similar domains	Pain reduction in 88% patients post-treatment with analgesics at $p < 0.001$ [58]	No reported floor-ceiling effects
Doloplus-2	Good inter-rater reliability of 0.73–0.81 [57] Test-retest reliability of 0.96 [26] High internal consistency of 0.85–0.86 at different time points [57]	Poor correlations with PAINAD, VAS and VRS with values ranging from 0.29–0.36 [26]	Significant reduction in mean scores after a given interval ($R^2 = 0.216$) [59]	No reported floor-ceiling effects
<i>Neuropathic pain-specific tool</i>				
NPS	Test intervals (about 33 days) produced test-retest reliability values of ICC = 0.71 as a total score, and individual ICC varied between 0.45–0.78 [60] Good internal consistency of 0.78 [60]	Excellent content validity with patients choosing to answer about 8 of the 10 items involved in this tool [60]; suggests that most articles in this tool are relevant to neuropathic pain One item allows patients to describe their pain in their own words	No reported responsiveness	No reported floor-ceiling effects

APS Abbey Pain Scale, *BPI* Brief Pain Inventory, *Doloplus* After the French investigator who developed the tool, *MPI* West Haven-Yale Multidimensional Pain Inventory, *MPQ* McGill Pain Questionnaire, *MVAS* Million’s Visual Analogue Scale, *MASS-LSO* The North American Spine Society Lumbar Spine Outcome, *NPS* Neuropathic Pain Scale, *ODI* Oswestry Disability Index, *PAINAD* Pain Assessment in Advanced Dementia, *PF* physical functioning, *PR* Physical role limitations, *BP* Bodily pain, *MH* Mental Health, *QBPD* Quebec Back Pain Disability Scale, *RMDQ* Roland Morris Disability Questionnaire, *SF-36* Short Form-36, *SRM* Standardized Response Mean, *WDJ* Waddell Disability Index

^aReliability measurement: 1. Inter-rater reliability: All values Pearson’s coefficient, intraclass coefficient (ICC) or Kappa coefficient. 2. Test-retest reliability: Values given as ICC or Pearson’s coefficient. 3. Internal consistency: Values given as Cronbach’s alpha (α)

^bConcurrent validity measurement: By comparison with other validated tools. All values Pearson’s correlation coefficient

^cResponsiveness measurement: Values given as effect size (ES; expressed as Pearson’s r or Cohen’s d), standardized response mean (SRM), or Receiver Operating Characteristic (ROC) curve. Area under this curve of 0.5 is poor and a value of 1.0 is perfect

^dFloor-ceiling effects: Values < 15% acceptable

^eThe statistical results show poor, adequate, good, and excellent values at < 0.5, 0.5–0.7, 0.7–0.8, > 0.9, respectively. The minimum acceptable limit for Cronbach’s α is 0.7, and a score > 0.9 indicates redundancy

Table 6 Merits and demerits of some major low back pain tools (generic and condition-specific)

Name of tool	Type of pain	Merits	Limitations	References
<i>Generic tools for back pain</i>				
BPI	Chronic/acute	Easy to score Can represent pain graphically Test–retest reliability high over short intervals High correlations with SF-36 and RMDQ Excellent responsiveness Cross-cultural use	Time-consuming	[28–30]
MPQ	Chronic/acute	Not unidimensional Allows collection of both qualitative and quantitative data Includes nociceptive and neuropathic items of sensory pain dimension Has good construct, content, and criterion validity Sensitive to intervention Strong reliability for cancer patients Psychometrically sound Cross-cultural use	Time-consuming Difficult to interpret Only three pain patterns not adequate for cancer patients	[16, 30, 31]
MPI	Chronic	Easy to administer Test–retest reliability across various domains ranges from adequate to excellent (depending upon language and disease) Construct validity adequate to excellent for chronic pain Psychometrically sound Cross-cultural use	Internal consistency scores disputed; ranges from adequate to excellent, or poor for certain domains like life control	[18, 33–35]
<i>HRQoL tool</i>				
SF-36	Chronic/acute	An HRQoL tool with an equally reliable and valid derivation, SF-12 Good to excellent reliability for PCS, excellent internal consistency for all domains Satisfactory correlations of physical functioning and pain with other disability tools Cross-cultural use	Poor reliability for mental health and social functioning Poor correlations of mental health with other disability tools Disputed responsiveness	[36–40, 61]
<i>Low back pain-specific tools</i>				
RMDQ-24	Chronic	Simple to use and fast completion Shows high correlation for physical function with other tools like ODI, SF-36, QBPDS Has good construct and concurrent criterion validity, internal consistency Test–retest reliability high over short intervals Cross-cultural use	Focusses on limited range of physical functions common only to patients with LBP; no option of “not applicable” on scoring system Poor test–retest reliability over long intervals Dichotomous response categories make it difficult to get much information Responsiveness depends on severity of disability Not very psychometrically sound	[41–43, 45]

Table 6 (continued)

Name of tool	Type of pain	Merits	Limitations	References
ODI	Chronic	Simple to use; fast completion and scoring Telephonic interview possible Shows good correlation with other tools like MPQ, VAS, SF-36 Has good construct validity, internal consistency Test–retest reliability high over short intervals Responsiveness acceptable for population of not severely affected people Good predictor of return-to-work Cross-cultural use	Poor test–retest reliability over long intervals Not so sound psychometrically	[39, 62, 63]
QBPDS	Chronic	Easy to administer and to score Possible in telephonic/mail interviews High test–retest reliability, good content and construct validity Has strong correlations with other disability tools Low item omission for activities asked to be performed Cross-cultural use	Has also been reported by some to be adequate to poor in terms of physical function and psychosocial measure Items on sex and social life not included	[19, 39, 40, 45]
MVAS	Chronic	Easy to administer and score; fast completion Mostly good test–retest reliability and internal consistency for daily activities and social life Has good construct validity and responsiveness Strong unidimensional measure of pain severity Cross-cultural use	Test–retest over various long and short intervals not tested Poor internal consistency for body functions Little psychometric evidence available	[50–52]
WDI	Chronic	Extremely simple and fast tool; easy to score Moderate test–retest reliability Good inter-rater reliability, construct validity Data can be pooled; advantage for clinical trials Cross-cultural use	Limited data on responsiveness	[9, 22, 40]
NASS-LSO	Chronic	High test–retest reliability over short intervals High internal consistency and construct validity Extensive low back pain-dominant questions Cross-cultural use	Extremely complex to administer and score Time-consuming No single questionnaire used; questions may be repeated Limited psychometric data	[23, 54]
<i>Tools for cognitively impaired</i>				
PAINAD	Chronic	Good construct validity and inter-rater and test–retest reliability Easy to administer and requires no intensive training as compared to other tools for patients with dementia Preferred tool of physicians Cross-cultural use	Not all actions presented by dementia patients may be caused by pain, for example, a sudden outburst may be due to emotional turmoil Internal consistency not acceptable, but can be improved by deleting “breathing” and “negative vocalization” Limited psychometric data	[24, 55]
APS	Chronic	Good internal consistency Good to excellent inter-rater reliability High correlations with PAINAD and DS-DAT Extremely good construct Takes less than a minute to complete Cross-cultural use	Construct of the tool not so sound; can be improved with removal of the item “physical change” Limited psychometric data	[56–58]

Table 6 (continued)

Name of tool	Type of pain	Merits	Limitations	References
Dolopius-2	Chronic	Simple to use Good inter-rater reliability and high internal consistency Excellent sensitivity to change Cross-cultural use	Takes 6–12 min to complete Construct validity only adequate Poor correlations with PAINAD, VAS and VRS Internal consistency and test–retest reliability controversial Requires extensive training	[26, 57, 59]
<i>Tool for neuropathic pain</i>				
NPS	Chronic	Good reliability and internal consistency Excellent content validity with high relevance to neuropathic pain Cross-cultural use	No data on responsiveness Limited psychometric data	[27, 60]
<i>Tool specifically for acute pain</i>				
CAPA	Acute	The only tool specific for acute pain Good patient and staff satisfaction	Psychometric data limited	[8, 64]

APS Abbey Pain Scale, *BPI* Brief Pain Inventory, *CAPA* Clinically Aligned Pain Assessment, *Dolopius* After the French investigator who developed the tool, *FPS* Faces Pain Scale, *MPI* West Haven-Yale Multidimensional Pain Inventory, *MPQ* McGill Pain Questionnaire, *MVAS* Million's Visual Analogue Scale, *NASS-LSO* The North American Spine Society Lumbar Spine Outcome, *NRS* Numerical Rating Scale, *ODI* Oswestry Disability Index, *PAINAD* Pain Assessment in Advanced Dementia, *NPS* Neuropathic Pain Scale, *QBPDS* Quebec Back Pain Disability Scale, *RMDQ* Roland Morris Disability Questionnaire, *SF-36* Short Form-36, *WDF* Waddell Disability Index, *VAS/GRS* Visual Analogue Scale/Graphic Rating Scale, *VRS* Verbal Rating Scale

disability, while ODI works well on people with severe disability of a persistent type [43]. ODI also has questions on a patient's sexual and social life providing a multifaceted pain assessment. QBPDS too is a functional tool like RMDQ and gauges only disability and sleep without measuring pain intensity [9]. It should be used in association with a pain-measuring unidimensional tool like NRS or VAS/GRS. WDI is a simple and short tool with an internal consistency higher than that of RMDQ but lower than ODI [40]. It has also been shown to detect significant changes in patients after a one-month interval post-treatment, but responsiveness has not been properly investigated [53]. This tool should be used along with other scoring systems for disability, to get a complete assessment of pain [9]. The inter-rater reliability of MVAS is not to be relied upon as there is little correlation between results of observers [21]. It has good internal consistency for all its subscales except body functions which is an important aspect for disability caused by LBP. The strength of MVAS lies in its VAS scale usage that provides a strong unidimensional measure of pain severity. The NASS-LSO is a tool with which one can easily mark the regions of pain on a picture provided to the patient. Though complex to use and score, and extremely time-consuming, this tool, if used, can provide a very comprehensive picture of pain. The strength of this tool lies in its questions focussed on low back pain [9, 23].

Among the generic measures reviewed here, all three measures, namely BPI, MPI, and MPQ, are strong, reliable, and valid tools for LBP. BPI and MPQ are comprehensive but time-consuming. BPI is easy to score, while MPI is easy to administer. Internal consistency of MPI ranges from adequate to excellent, depending upon the language version used and the disease being studied. Construct validity of the tool too ranges from acceptable to excellent proving that all items being investigated serve the purpose of the tool. All three tools have cross-cultural use and show good responsiveness, with MPQ being extremely sensitive to detect change due to intervention. BPI shows high test–retest ability [28]. MPQ has the added advantage of including both nociceptive and neuropathic items of sensory pain dimension and is thus used for neuropathic LBP too. MPQ, MPI, and BPI are all used to assess diverse chronic pain conditions, including LBP, but BPI is the only tool that is formally validated for LBP, although not chronic LBP [66]. However, certain problems may arise in using BPI as patients' answers rely more on the disability caused by LBP in the recent past as opposed to the pain suffered, in general, during daily activities. In addition to this, when asked to fill the questionnaire, there may be a bias (patient-related) based on the patient's ability to bear the pain. The former can be overcome by noting the findings, in accordance with the questionnaire, after hearing the chief complaints of the patient. This prevents wastage of time and redundancy in the clinical queries. The latter can

Table 7 Mnemonic tools for quick assessment of acute pain

S. no.	Name of tool	Description	References
1.	SOCRATES	Site: Pain location? Onset: When did pain start and how quickly did it progress? Character: Dull, stabbing or sharp pain? Radiation: Does the pain spread to other body parts? Associations: Is pain present with other factors like nausea or movement difficulty, fever? Time: Is any pattern followed by the pain? Exacerbating/relieving factors: is pain worse at rest, movement or with certain basic functions like breathing? Surgical history: Any previous surgery or interventions?	[8]
2.	PQRST	<i>P</i> Precipitating factors <i>Q</i> Quality of pain <i>R</i> Radiating pain <i>S</i> Severity of pain <i>T</i> Temporal factor	[8]
3.	QISS-TAPED	<i>Q</i> Quality <i>I</i> Impact <i>S</i> Site <i>S</i> Severity <i>T</i> Temporal characteristics <i>A</i> Aggravating and alleviating factors <i>P</i> Past treatment, response and patient preferences <i>E</i> Expectations and meaning <i>D</i> Diagnostics and physical examination	[8]

be solved by blinding the patient to the assessment process which helps in getting more honest answers.

SF-36 is a very strong tool for the assessment of general health-related quality of life and also as an LBP tool. The tool's bodily pain and physical function correlate well with other disability tools, but the mental health subscale results are poor [38, 39].

Among the tools for assessing neuropathic pain, NPS seems like the only generic tool, apart from SF-MPQ-2, that has adequate validation and is widely accepted [27, 60]. This tool has good internal consistency and test–retest values as proven in a study by Rog et al. [60]. Though no formal responsiveness has been reported for this tool, and more confirmatory work is required on its psychometric properties, this tool has proven to be useful for assessment of neuropathic pain [27, 60].

Even in patients with dementia, a self-reported pain tool should always be the first attempt in order to estimate pain severity along with determining the presence or absence of pain, but due to the inability to execute the generic and condition-specific tools among these subjects, we must rely on observational measures. Among the tools for assessment of LBP among elderly or people who are cognitively impaired, all three tools have proven to be reliable, but no formal responsiveness has been tested for the tools. PAINAD shows excellent inter-rater reliabilities during pleasant and unpleasant activities [24] but poor to moderate reliability at rest or during movement [55]. It is thus very good at discriminating between different conditions like rest, movement, or pleasant

and unpleasant activities, establishing a good construct validity [24, 55]. It shows good correlations with the pain and discomfort scale of VAS, and with DS-DAT. Internal consistency is poor and does not achieve the accepted limit of 0.70 Cronbach's α , but removing the item "breathing" or "negative vocalization" leads to an acceptable internal consistency of 0.75 [24]. Doloplus-2 shows high internal consistency and inter-rater reliability [57]. Test–retest values are extremely good, but this tool does not correlate well with other tools like PAINAD, VAS, and VRS [26]. APS shows good to excellent values for both inter-rater reliability and internal consistency, and removing the item "physical change" improves the fit of the tool, and thus its construct validity [56]. Both APS and Doloplus-2 show high sensitivity to change after a certain interval or intervention [58]. For acute pain assessment, unidimensional tools seem to be the norm, though tools like BPI and MPQ are also used considerably. CAPA, even with its limited psychometric data, has proven to be reliable, responsive, and most importantly, consistent in providing patient and staff satisfaction, as evident from the results of University of Utah Hospital [8, 64]. This tool needs to be tested more and across continents to attain a better understanding of its quality.

Conclusion

Every tool reviewed here has its own advantages and limitations. Recommendations would vary depending upon the language, disease condition, pain severity, and most of all the physician and patient using the tool. However, among the LBP-specific tools, RMDQ and ODI seem to be the ones most widely accepted and can help in clinical conditions like prolapsed intervertebral disc, infective spine, spondylosis, and spondylolisthesis. They need to be used in conjunction with a HRQoL tool like SF-36 or its derived version SF-12, but SF-36 can be used as an independent measure for LBP too. Among the generic tools for LBP, even though both BPI and MPQ are time-consuming, they have high reliability and validity. Our review points to a higher level of proven responsiveness for BPI as compared to MPQ. Also, BPI has been formally validated for LBP, making it the best generic tool for LBP management at present, in our opinion. For neuropathic LBP, more intervention studies are required to acquire data on the sensitivity of NPS. In the meantime, it can be used together with SF-MPQ-2 to get an accurate idea of the pain encountered. These neuropathic pain tools can evaluate neurological deficits which in turn can help to treat the patients conservatively or surgically. PAINAD, APS, and Doloplus-2 are all acceptable tools for cognitively impaired people and the choice of using one or more rests on the physicians. We understand that unidimensional tools, SF-MPQ-2 and CAPA, can all be used successfully for acute low back pain. Even the non-validated quick assessment mnemonic tools may not be irrelevant, as for acute pain, quick assessment and quick relief is the fundamental goal. Lack of formal validation of a pain tool has not stopped its worldwide use and acceptability in daily clinical practice. Rather, these mnemonic tools can help in the clinical evaluation of acute pain in orthopaedic diseases like prolapsed intervertebral disc, spinal canal stenosis, spondylosis and diseases like infective spine and ankylosing spondylitis. Finally, physicians and patients need to use the tool that is best suited for the specific condition and pain severity.

Acknowledgements We thank Turacoz Healthcare Solutions (www.turacoz.com), Gurugram, India for writing support.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Human and animal rights This article is based on previously conducted studies and does not contain any studies with human participants or animals performed by any of the authors.

References

1. Finger ME, Selb M, De Bie R, Escorpizo R (2015) Using the international classification of functioning, disability and health in physiotherapy in multidisciplinary vocational rehabilitation: a case study of low back pain. *Physiother Res Int* 20(4):231–241. <https://doi.org/10.1002/pri.1587>
2. Freburger JK, Holmes GM, Agans RP, Jackman AM, Darter JD, Wallace AS, Castel LD, Kalsbeek WD, Carey TS (2009) The rising prevalence of chronic low back pain. *Arch Intern Med* 169(3):251–258. <https://doi.org/10.1001/archinternmed.2008.543>
3. Hartvigsen J, Hancock MJ, Kongsted A, Louw Q, Ferreira ML, Genevay S, Hoy D, Karppinen J, Pransky G, Sieper J, Smeets RJ, Underwood M (2018) What low back pain is and why we need to pay attention. *Lancet* 391(10137):2356–2367. [https://doi.org/10.1016/s0140-6736\(18\)30480-x](https://doi.org/10.1016/s0140-6736(18)30480-x)
4. Atlas SJ, Chang Y, Kammann E, Keller RB, Deyo RA, Singer DE (2000) Long-term disability and return to work among patients who have a herniated lumbar disc: the effect of disability compensation. *J Bone Joint Surg Am* 82(1):4–15
5. Hoy D, March L, Brooks P, Blyth F, Woolf A, Bain C, Williams G, Smith E, Vos T, Barendregt J, Murray C, Burstein R, Buchbinder R (2014) The global burden of low back pain: estimates from the Global Burden of Disease 2010 study. *Ann Rheum Dis* 73(6):968–974. <https://doi.org/10.1136/annrheumdis-2013-204428>
6. Jay K, Friberg MK, Sjogaard G, Jakobsen MD, Sundstrup E, Brandt M, Andersen LL (2015) The consequence of combined pain and stress on work ability in female laboratory technicians: a cross-sectional study. *Int J Environ Res Public Health* 12(12):15834–15842. <https://doi.org/10.3390/ijerph121215024>
7. Caumo W, Ruehlman LS, Karoly P, Sehn F, Vidor LP, Dall-Agnol L, Chassot M, Torres IL (2013) Cross-cultural adaptation and validation of the profile of chronic pain: screen for a Brazilian population. *Pain Med* 14(1):52–61. <https://doi.org/10.1111/j.1526-4637.2012.01528.x>
8. Gordon DB (2015) Acute pain assessment tools: let us move beyond simple pain ratings. *Curr Opin Anaesthesiol* 28(5):565–569. <https://doi.org/10.1097/aco.0000000000000225>
9. Longo UG, Loppini M, Denaro L, Maffulli N, Denaro V (2010) Rating scales for low back pain. *Br Med Bull* 94:81–144. <https://doi.org/10.1093/bmb/ldp052>
10. Muller U, Duetz MS, Roeder C, Greenough CG (2004) Condition-specific outcome measures for low back pain. Part I: validation. *Eur Spine J* 13(4):301–313. <https://doi.org/10.1007/s00586-003-0665-1>
11. Turk DC, Dworkin RH, Allen RR, Bellamy N, Brandenburg N, Carr DB, Cleeland C, Dionne R, Farrar JT, Galer BS, Hewitt DJ, Jadad AR, Katz NP, Kramer LD, Manning DC, McCormick CG, McDermott MP, McGrath P, Quessy S, Rappaport BA, Robinson JP, Royal MA, Simon L, Stauffer JW, Stein W, Tollett J, Witter J (2003) Core outcome domains for chronic pain clinical trials: IMMPACT recommendations. *Pain* 106(3):337–345
12. Turk DC, Dworkin RH, Revicki D, Harding G, Burke LB, Cella D, Cleeland CS, Cowan P, Farrar JT, Hertz S, Max MB, Rappaport BA (2008) Identifying important outcome domains for chronic pain clinical trials: an IMMPACT survey of people with pain. *Pain* 137(2):276–285. <https://doi.org/10.1016/j.pain.2007.09.002>
13. Cleeland CS, Ryan KM (1994) Pain assessment: global use of the brief pain inventory. *Ann Acad Med Singapore* 23(2):129–138
14. Roland M, Morris R (1983) A study of the natural history of back pain. Part I: development of a reliable and sensitive measure of disability in low-back pain. *Spine (Phila Pa 1976)* 8(2):141–144

15. Melzack R, Katz J (2001) The McGill pain questionnaire: appraisal and current status. Handbook of pain assessment, 2nd edn. The Guilford Press, New York, pp 35–52
16. Dworkin RH, Turk DC, Revicki DA, Harding G, Coyne KS, Peirce-Sandner S, Bhagwat D, Everton D, Burke LB, Cowan P, Farrar JT, Hertz S, Max MB, Rappaport BA, Melzack R (2009) Development and initial validation of an expanded and revised version of the Short-form McGill Pain Questionnaire (SF-MPQ-2). *Pain* 144(1–2):35–42. <https://doi.org/10.1016/j.pain.2009.02.007>
17. Fairbank JC, Pynsent PB (2000) The Oswestry disability index. *Spine (Phila Pa 1976)* 25(22):2940–2952 **discussion 2952**
18. Kerns RD, Turk DC, Rudy TE (1985) The west haven-yale multidimensional pain inventory (WHYMPI). *Pain* 23(4):345–356
19. Kopec JA, Esdaile JM, Abrahamowicz M, Abenhaim L, Wood-Dauphinee S, Lamping DL, Williams JI (1995) The Quebec Back Pain Disability Scale. Measurement properties. *Spine (Phila Pa 1976)* 20(3):341–352
20. Ware JE Jr, Sherbourne CD (1992) The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 30(6):473–483
21. Million R, Hall W, Nilsen KH, Baker RD, Jayson MI (1982) Assessment of the progress of the back-pain patient 1981 Volvo Award in Clinical Science. *Spine (Phila Pa 1976)* 7(3):204–212
22. Waddell G, Main CJ (1984) Assessment of severity in low-back disorders. *Spine (Phila Pa 1976)* 9(2):204–208
23. Daltroy LH, Cats-Baril WL, Katz JN, Fossel AH, Liang MH (1996) The North American spine society lumbar spine outcome assessment Instrument: reliability and validity tests. *Spine (Phila Pa 1976)* 21(6):741–749
24. Warden V, Hurley AC, Volicer L (2003) Development and psychometric evaluation of the pain assessment in advanced dementia (PAINAD) scale. *J Am Med Dir Assoc* 4(1):9–15. <https://doi.org/10.1097/01.Jam.0000043422.31640.F7>
25. Abbey J, Pillar N, De Bellis A, Esterman A, Parker D, Giles L, Lowcay B (2004) The Abbey pain scale: a 1-minute numerical indicator for people with end-stage dementia. *Int J Palliat Nurs* 10(1):6–13. <https://doi.org/10.12968/ijpn.2004.10.1.12013>
26. Pautex S, Herrmann FR, Michon A, Giannakopoulos P, Gold G (2007) Psychometric properties of the Doloplus-2 observational pain assessment scale and comparison to self-assessment in hospitalized elderly. *Clin J Pain* 23(9):774–779. <https://doi.org/10.1097/AJP.0b013e318154b6e3>
27. Galer BS, Jensen MP (1997) Development and preliminary validation of a pain measure specific to neuropathic pain: the neuropathic pain scale. *Neurology* 48(2):332–338
28. Erdemoglu AK, Koc R (2013) Brief Pain Inventory score identifying and discriminating neuropathic and nociceptive pain. *Acta Neurol Scand* 128(5):351–358. <https://doi.org/10.1111/ane.12131>
29. Keller S, Bann CM, Dodd SL, Schein J, Mendoza TR, Cleeland CS (2004) Validity of the brief pain inventory for use in documenting the outcomes of patients with noncancer pain. *Clin J Pain* 20(5):309–318
30. Dworkin RH, Turk DC, Trudeau JJ, Benson C, Biondi DM, Katz NP, Kim M (2015) Validation of the short-form McGill Pain Questionnaire-2 (SF-MPQ-2) in acute low back pain. *J Pain* 16(4):357–366. <https://doi.org/10.1016/j.jpain.2015.01.012>
31. Kachooei AR, Ebrahimzadeh MH, Erfani-Sayyar R, Salehi M, Salimi E, Razi S (2015) Short Form-McGill Pain Questionnaire-2 (SF-MPQ-2): a cross-cultural adaptation and validation study of the persian version in patients with knee osteoarthritis. *Arch Bone Jt Surg* 3(1):45–50
32. Strand LI, Ljunggren AE, Bogen B, Ask T, Johnsen TB (2008) The Short-Form McGill Pain Questionnaire as an outcome measure: test–retest reliability and responsiveness to change. *Eur J Pain* 12(7):917–925. <https://doi.org/10.1016/j.ejpain.2007.12.013>
33. Verra ML, Angst F, Staal JB, Brioschi R, Lehmann S, Aeschlimann A, de Bie RA (2012) Reliability of the Multidimensional Pain Inventory and stability of the MPI classification system in chronic back pain. *BMC Musculoskelet Disord* 13:155. <https://doi.org/10.1186/1471-2474-13-155>
34. Angst F, Verra ML, Lehmann S, Aeschlimann A (2008) Responsiveness of five condition-specific and generic outcome assessment instruments for chronic pain. *BMC Med Res Methodol* 8(1):26. <https://doi.org/10.1186/1471-2288-8-26>
35. Cetin AA, Bektas H, Ozdogan M (2016) The west haven yale multidimensional pain inventory: reliability and validity of the Turkish version in individuals with cancer. *Eur J Oncol Nurs* 20:1–9. <https://doi.org/10.1016/j.ejon.2015.03.007>
36. Busija L, Pausenberger E, Haines TP, Haymes S, Buchbinder R, Osborne RH (2011) Adult measures of general health and health-related quality of life: Medical Outcomes Study Short Form 36-Item (SF-36) and Short Form 12-Item (SF-12) Health Surveys, Nottingham Health Profile (NHP), Sickness Impact Profile (SIP), Medical Outcomes Study Short Form 6D (SF-6D), Health Utilities Index Mark 3 (HUI3), Quality of Well-Being Scale (QWB), and Assessment of Quality of Life (AQL). *Arthritis Care Res (Hoboken)* 63(Suppl 11):S383–412. <https://doi.org/10.1002/acr.20541>
37. White MK, McCausland KL, Sancherawala V, Guthrie SD, Bayliss MS (2017) Psychometric validation of the SF-36 Health Survey in light chain amyloidosis: results from community-based and clinic-based samples. *Patient Relat Outcome Meas* 8:157–167. <https://doi.org/10.2147/prom.S146849>
38. Chiarotto A, Maxwell LJ, Terwee CB, Wells GA, Tugwell P, Ostelo RW (2016) Roland–Morris Disability Questionnaire and Oswestry Disability Index: which has better measurement properties for measuring physical functioning in nonspecific low back pain? systematic review and meta-analysis. *Phys Ther* 96(10):1620–1637. <https://doi.org/10.2522/ptj.20150420>
39. Hicks GE, Manal TJ (2009) Psychometric properties of commonly used low back disability questionnaires: are they useful for older adults with low back pain? *Pain Med* 10(1):85–94. <https://doi.org/10.1111/j.1526-4637.2008.00548.x>
40. Davidson M, Keating JL (2002) A comparison of five low back disability questionnaires: reliability and responsiveness. *Phys Ther* 82(1):8–24
41. Stratford PW, Binkley J, Solomon P, Finch E, Gill C, Moreland J (1996) Defining the minimum level of detectable change for the Roland–Morris questionnaire. *Phys Ther* 76(4):359–365 (**discussion 366–358**)
42. Payares K, Lugo LH, Restrepo A (2015) Validation of the Roland Morris Questionnaire in Colombia to evaluate disability in low back pain. *Spine (Phila Pa 1976)* 40(14):1108–1114. <https://doi.org/10.1097/brs.0000000000000963>
43. Roland M, Fairbank J (2000) The Roland–Morris disability questionnaire and the Oswestry disability questionnaire. *Spine (Phila Pa 1976)* 25(24):3115–3124
44. Kopec JA, Esdaile JM (1995) Functional disability scales for back pain. *Spine (Phila Pa 1976)* 20(17):1943–1949
45. Reneman MF, Jorritsma W, Schellekens JM, Goeken LN (2002) Concurrent validity of questionnaire and performance-based disability measurements in patients with chronic nonspecific low back pain. *J Occup Rehabil* 12(3):119–129
46. Monticone M, Baiardi P, Vanti C, Ferrari S, Pillastrini P, Mugnai R, Foti C (2012) Responsiveness of the Oswestry Disability Index and the Roland Morris Disability Questionnaire in Italian subjects with sub-acute and chronic low back pain. *Eur Spine J* 21(1):122–129. <https://doi.org/10.1007/s00586-011-1959-3>
47. Müller U, Röder C, Greenough CG (2006) Back related outcome assessment instruments. *Eur Spine J* 15(Suppl 1):S25–S31. <https://doi.org/10.1007/s00586-005-1054-8>

48. Fairbank JC, Couper J, Davies JB, O'Brien JP (1980) The Oswestry low back pain disability questionnaire. *Physiotherapy* 66(8):271–273
49. Payares K, Lugo LH, Morales V, Londono A (2011) Validation in Colombia of the Oswestry disability questionnaire in patients with low back pain. *Spine* 36(26):E1730–E1735. <https://doi.org/10.1097/brs.0b013e318219d184>
50. Akbari H, Ghasemi M, Yegani T, Fesharaki MG, Saraei M, Barsam Y, Akbari H (2018) Million Visual Analogue Scale Questionnaire: validation of the Persian Version. *Asian Spine J.* <https://doi.org/10.31616/asj.2018.0223>
51. Alaranta H, Rytokoski U, Rissanen A, Talo S, Ronnema T, Puukka P, Karppi SL, Videman T, Kallio V, Slati P (1994) Intensive physical and psychosocial training program for patients with chronic low back pain. A controlled clinical trial. *Spine (Phila Pa 1976)* 19(12):1339–1349
52. Anagnostis C, Mayer TG, Gatchel RJ, Proctor TJ (2003) The million visual analog scale: its utility for predicting tertiary rehabilitation outcomes. *Spine (Phila Pa 1976)* 28(10):1051–1060. <https://doi.org/10.1097/01.brs.0000061989.94487.9b>
53. Ruta DA, Garratt AM, Wardlaw D, Russell IT (1994) Developing a valid and reliable measure of health outcome for patients with low back pain. *Spine (Phila Pa 1976)* 19(17):1887–1896
54. Sarasqueta C, Gabaldon O, Iza I, B eland F, Paz PM (2005) Cross-cultural adaptation and validation of the NASS outcomes instrument in Spanish patients with low back pain. *Eur Spine J* 14(6):586–594. <https://doi.org/10.1007/s00586-004-0871-5>
55. Ersek M, Herr K, Neradilek MB, Buck HG, Black B (2010) Comparing the psychometric properties of the checklist of nonverbal pain behaviors (CNPI) and the pain assessment in advanced dementia (PAIN-AD) instruments. *Pain Med* 11(3):395–404. <https://doi.org/10.1111/j.1526-4637.2009.00787.x>
56. Liu JY, Briggs M, Closs SJ (2010) The psychometric qualities of four observational pain tools (OPTs) for the assessment of pain in elderly people with osteoarthritic pain. *J Pain Symptom Manag* 40(4):582–598. <https://doi.org/10.1016/j.jpainsymman.2010.02.022>
57. Neville C, Ostini R (2014) A psychometric evaluation of three pain rating scales for people with moderate to severe dementia. *Pain Manag Nurs* 15(4):798–806. <https://doi.org/10.1016/j.pmn.2013.08.001>
58. Gregersen M, Melin AS, Nygaard IS, Nielsen CH, Beedholm-Ebsen M (2016) Reliability of the Danish Abbey Pain Scale in severely demented and non-communicative older patients. *Int J Palliat Nurs* 22(10):482–488. <https://doi.org/10.12968/ijpn.2016.22.10.482>
59. Monacelli F, Vasile Nurse A, Odetti P, Traverso N (2013) Doloplus-2 pain assessment: an effective tool in patients over 85 years with advanced dementia and persistent pain. *Clin Ter* 164(1):e23–25. <https://doi.org/10.7417/ct.2013.1516>
60. Rog DJ, Nurmikko TJ, Friede T, Young CA (2007) Validation and reliability of the neuropathic pain scale (NPS) in multiple sclerosis. *Clin J Pain* 23(6):473–481. <https://doi.org/10.1097/AJP.0b013e31805d0c5d>
61. Riddle DL, Lee KT, Stratford PW (2001) Use of SF-36 and SF-12 health status measures: a quantitative comparison for groups versus individual patients. *Med Care* 39(8):867–878
62. Hupli M, Heinonen R, Vanharanta H (1997) Height changes among chronic low back pain patients during intense physical exercise. *Scand J Med Sci Sports* 7(1):32–37
63. Nordin M, Skovron ML, Hiebert R, Weiser S, Brisson PM, Campello M, Harwood K, Crane M, Lewis S (1997) Early predictors of delayed return to work in patients with low back pain. *J Musculoskelet Pain* 5(2):5–27
64. Topham D, Drew D (2017) Quality improvement project: replacing the numeric rating scale with a clinically aligned pain assessment (CAPA) tool. *Pain Manag Nurs* 18(6):363–371. <https://doi.org/10.1016/j.pmn.2017.07.001>
65. Chapman JR, Norvell DC, Hermsmeyer JT, Bransford RJ, DeVine J, McGirt MJ, Lee MJ (2011) Evaluating common outcomes for measuring treatment success for chronic low back pain. *Spine (Phila Pa 1976)* 36(21 Suppl):S54–S68. <https://doi.org/10.1097/brs.0b013e31822ef74d>
66. Ramasamy A, Martin ML, Blum SI, Liedgens H, Argoff C, Freynhagen R, Wallace M, McCarrier KP, Bushnell DM, Hatley NV, Patrick DL (2017) Assessment of patient-reported outcome instruments to assess chronic low back pain. *Pain Med* 18(6):1098–1110. <https://doi.org/10.1093/pm/pnw357>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.